

Quantum optical implementation of Grover's algorithm

Marlan O. Scully*†‡ and M. S. Zubairy*§

*Department of Physics, Texas A&M University, College Station, TX 77843-4242; †Max-Planck-Institut für Quantenoptik, 85748 Garching, Germany; and §Department of Electronics, Quaid-i-Azam University, Islamabad, Pakistan

Contributed by Marlan O. Scully, June 21, 2001

We present a scheme for a quantum optical implementation of Grover's algorithm based on resonant atomic interactions with classical fields and dispersive couplings with quantized cavity fields. The proposed scheme depends on preparation of entangled states and is within current state-of-the-art technology.

As was first shown by Grover (1), search of a database by using quantum mechanics can be substantially faster than any classical search of unsorted data. For example, it was shown by Grover that, by using quantum superpositions and quantum entanglement, we can find an object in an unsorted database containing N objects in $O(\sqrt{N})$ quantum mechanical steps instead of $O(N)$ steps (1–3).

The implementation of search algorithms by using optical methods is a subject of intense interest (4–8), and Grover's search algorithm has been implemented by using nuclear magnetic resonance (NMR) techniques for a system with four states (9–11). However, NMR experiments for quantum computing are carried out at room temperature, and questions have been raised concerning the appearance of entanglement in the physical state at any stage of such experiments (12). Braunstein *et al.* show that “all states so far used in NMR for quantum computations or for other quantum-information protocols are separable,” and therefore “no entanglement appears in the physical states at any stage of present NMR experiments.”

In this paper, we propose a scheme for quantum optical implementation of Grover's algorithm that is not sensitive to such thermal decoherence effects. The scheme is based on resonant atomic interactions with classical fields and dispersive coupling with quantized cavity fields. We first formulate the problem in terms of a “circuit” logic involving one-bit unitary transformation and a two-bit quantum phase gate. For an atomic system, the one-bit unitary transformation is accomplished by means of resonant interaction with a classical field, whereas a quantum phase gate can be implemented by using dispersive coupling with a cavity field having either 0 or 1 photon. Such a quantum phase gate has been demonstrated recently [for a beautiful demonstration of quantum phase gate, see Rauschenbeutel *et al.* (13); conditional phase shifts are also demonstrated in Turchette *et al.* (14)]. The proposed scheme involving atomic interaction with classical field and two cavities therefore lies within present experimental limitations and should be realizable (15).

Grover proposed an algorithm to search an item in an unsorted database. The problem he addressed is as follows. We are given a function $f(x)$ with $x = 1, 2, \dots, N$. The function has the property that it is 0 for all values of x except for x_0 , for which $f(x_0) = 1$. The task is to find x_0 . Classically, it would require an average of $N/2$ steps to accomplish this task. Grover showed that we can find x_0 in $O(\sqrt{N})$ quantum mechanical steps instead of $O(N)$ steps.

The basic idea of Grover's algorithm is to invert the phase (e.g., change $+$ → $-$, as in the passage from Eqs. 6 to 8) of the desired basis state and then invert all the basis states about the average amplitude of all the states. In this paper, we restrict ourselves to the simplest interesting Grover's algorithm with $N =$

4 with two qubits. First, we discuss the implementation of Grover's algorithm in terms of quantum logic gates.

A universal quantum computer consists of only two gates, namely a unitary transformation (one-bit gate) and a two-bit conditional quantum phase gate. The one-bit quantum gate for the i th qubit is given by

$$U_{\theta,\phi}^i = \begin{pmatrix} \cos\theta & -ie^{-i\phi}\sin\theta \\ -ie^{i\phi}\sin\theta & \cos\theta \end{pmatrix}. \quad [1]$$

A convenient representation of $U_{\theta,\phi}$ in terms of Pauli spin matrices is given by

$$U_{\theta,\phi} = \cos\theta I - i\cos\phi\sin\theta\sigma_x - i\sin\phi\sin\theta\sigma_y. \quad [2]$$

The transformation for a two-bit quantum phase gate is given by $Q_\eta|\alpha, \beta\rangle = \exp(i\eta\delta_{\alpha,1}\delta_{\beta,1})|\alpha, \beta\rangle$, where $|\alpha\rangle$ and $|\beta\rangle$ stand for the basis states $|0\rangle$ or $|1\rangle$ of the qubit. Thus, the quantum phase gate introduces a phase η only when both qubits in the input state are 1. In the following, we shall need the quantum phase gate only with $\eta = \pi$, for which we have

$$Q_\pi = |0, 0\rangle\langle 0, 0| + |0, 1\rangle\langle 0, 1| + |1, 0\rangle\langle 1, 0| - |1, 1\rangle\langle 1, 1|, \quad [3]$$

and because $|0\rangle\langle 0| = (1 + \sigma_z)/2$ and $|1\rangle\langle 1| = (1 - \sigma_z)/2$,

$$Q_\pi = \frac{1}{2}(1_1 1_2 + 1_1 \sigma_{z2} + \sigma_{z1} 1_2 - \sigma_{z1} \sigma_{z2}). \quad [4]$$

Grover's algorithm is accomplished in three steps. The initial state of the two qubits is $|1, 1\rangle$. In the first step, we apply the Walsh–Hadamard transformation,

$$\mathcal{W} = \frac{(1 + i\sigma_{y1})}{\sqrt{2}} \frac{(1 + i\sigma_{y2})}{\sqrt{2}} = U_{\pi/4, -\pi/2}^1 U_{\pi/4, -\pi/2}^2, \quad [5]$$

which rotates each qubit from $|0\rangle$ to $(|0\rangle - |1\rangle)/\sqrt{2}$ and $|1\rangle$ to $(|0\rangle + |1\rangle)/\sqrt{2}$. The resulting state is

$$|s\rangle = \frac{1}{2}(|0, 0\rangle + |0, 1\rangle + |1, 0\rangle + |1, 1\rangle). \quad [6]$$

In the second step, the unitary operator $C_{\alpha,\beta}$ flips the sign of state $|\alpha, \beta\rangle$ ($\alpha = 0$ or 1 and $\beta = 0$ or 1). In the original Grover's algorithm, this is accomplished through operator $(1 - 2|\alpha, \beta\rangle\langle \alpha, \beta|)$. Here we follow a different approach. We first flip the sign of state $|1, 1\rangle$ via a quantum phase gate Q_π , followed by unitary operators that either retain the state of the qubit or change the state of the qubit from 0 to 1 and 1 to 0. The sign flip operators for the four possible states in this approach are given by

$$C_{0,0} = -\sigma_{x1}\sigma_{x2}Q_\pi = U_{\pi/2,0}^1 U_{\pi/2,0}^2 Q_\pi, \quad [7a]$$

*To whom reprint requests may be addressed. E-mail: scully@physics.tamu.edu or zubairy@physics.tamu.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. §1734 solely to indicate this fact.




Fig. 1. Level diagram for the implementation of Grover's algorithm.

$$C_{0,1} = -i\sigma_{x1}1_2Q_\pi = U_{\pi/2,0}^1Q_\pi, \quad [7b]$$

$$C_{1,0} = -i1_1\sigma_{x2}Q_\pi = U_{\pi/2,0}^2Q_\pi, \quad [7c]$$

$$C_{1,1} = Q_\pi. \quad [7d]$$

Here the resulting state may have an unimportant overall phase factor.

At this point, the oracle applies one of the $C_{\alpha,\beta}$ operators to state 6 and thus changes one of the signs from + to -. For example, if $C_{0,1}$ is chosen, then Eq. 6 becomes

$$C_{0,1}|s\rangle = \frac{-i}{2}(|0,0\rangle - |0,1\rangle + |1,0\rangle + |1,1\rangle). \quad [8]$$

It is now our job to find the marked state ($|0,1\rangle$ in the above example). This is accomplished by the application of operator $\mathcal{N} = 1 - 2|s\rangle\langle s|$. The resulting state is the desired state $|a, b\rangle$ apart from an unimportant π phase shift. To find a representation for \mathcal{N} in terms of operators $U_{\theta,\phi}$ and Q_π , we first note that

$$\mathcal{N} = \frac{1}{2}(1_11_2 - 1_1\sigma_{x2} - \sigma_{x1}1_2 - \sigma_{x1}\sigma_{x2}), \quad [9]$$

so that, from Eqs. 8 and 9, we have $\mathcal{N}C_{0,1}|s\rangle = -|0,1\rangle$. If we use the fact that rotating σ_z about the y axis yields σ_x , that is,

$$\frac{(1+i\sigma_y)}{\sqrt{2}}\sigma_z\frac{(1-i\sigma_y)}{\sqrt{2}} = -\sigma_x, \quad [10]$$

we note the remarkable fact that operator \mathcal{N} as given by Eq. 9 is the appropriately rotated quantum phase operator Q_π of Eq. 7b, i.e.,

$$\mathcal{N} = U_{\pi/4, -\pi/2}^1U_{\pi/4, -\pi/2}^2Q_\piU_{\pi/4, \pi/2}^1U_{\pi/4, \pi/2}^2. \quad [11]$$

This is the essence of Grover's algorithm and the departure point for our implementation of the algorithm.

On combining the various operators from Eqs. 5, 7, and 11, we can write Grover's algorithm in terms of simple transformations corresponding to one- and two-bit quantum gates as $\mathcal{N}C_{\alpha,\beta}\mathcal{W}|1, 1\rangle = e^{i\psi}|\alpha, \beta\rangle$, where ψ is a phase depending on the choice of α and β . These steps are summarized in the “circuit” diagram of Fig. 1 for the present ($N = 4$) implementation of Grover's algorithm. Only the one-bit unitary gates and two-bit conditional quantum phase gates are required for this purpose.

Next, we consider schemes for the implementation of Grover's algorithm on the basis of cavity quantum electrodynamics (16–20), using resonant atomic interaction with classical Ramsey fields and dispersive atomic interaction with quantized high Q cavity fields. First, we present a scheme that requires two species of atoms and two cavities. This scheme is conceptually simple but difficult to implement. We then consider a second scheme that

requires single species of atoms interacting with only one cavity that supports two modes, and the atomic level spacings are appropriately changed by electric or magnetic fields.

First, we consider the schematic for the proposed implementation as given in Fig. 2. The two qubits are represented by two different three-level atoms of types A and B (Fig. 2a). Levels $|a\rangle$ and $|b\rangle$ of these atoms represent qubits $|0\rangle$ and $|1\rangle$, respectively. The levels scheme is such that $\omega_{a_1}b_1 = \omega_{c_1}b_2 + \Delta_1$ and $\omega_{a_2}b_2 = \omega_{c_1}b_1 + \Delta_2$. Cavities C_1 and C_2 are resonant with transitions $\omega_{a_1}b_1$ and $\omega_{a_2}b_2$, respectively. Here, and in the following, the odd subscripts 1, 3, 5, and 7 refer to the atom, cavity, and classical fields corresponding to the atoms of type A, whereas the even subscripts 2, 4, 6, and 8 refer to atoms of type B. Four atoms (two of type A and two of type B) pass through cavities C_1 and C_2 and a sequence of classical fields, as follows (Fig. 2b).

All the atoms are initially in their ground states $|b\rangle$, i.e., $|b_1, b_2, b_3, b_4\rangle$, and we assume that the classical Ramsey fields are on only when the appropriate atoms are passing through them. Atom 1 of type A and atom 2 of type B are initially in their ground states $|b_1\rangle$ and $|b_2\rangle$, i.e., in qubits $|1, 1\rangle$. The Walsh–Hadamard transformation on these atoms is carried out by interacting with classical fields R_1 and R_2 of frequencies $\omega_{a_1}b_1$ and $\omega_{a_2}b_2$, respectively. The interaction of an atom with the classical field results in the unitary transformation (Eq. 1) on the atomic states $|a\rangle$ and $|b\rangle$, such that θ depends on the Rabi frequency and the interaction time, and ϕ depends on the phase of the driving field (21, 22). We choose, for the interaction of R_1 with atom 1 and R_2 with atom 2, $\theta_1 = \theta_2 = \pi/4$ and $\phi_1 = \phi_2 = -\pi/2$, so that the state of the atoms is $(|a_1, a_2\rangle + |a_1, b_2\rangle + |b_1, a_2\rangle + |b_1, b_2\rangle)/\sqrt{2}$.

In the next step, we consider operation $C_{\alpha,\beta}$. This step requires, in addition to the unitary operations, a quantum phase gate. The operation of the quantum phase gate is accomplished as follows. Atom 1 in the state $(|a_1\rangle + |b_1\rangle)/\sqrt{2}$ passes through the empty cavity C_1 . The interaction time is chosen such that the atom leaves in the ground state $|b_1\rangle$, and the field state inside the cavity becomes $(|0_1\rangle + |1_1\rangle)/\sqrt{2}$. Atom 2, which is of type B, then passes through the cavity. Because of the dispersive coupling, the effective Hamiltonian for the interaction between atom 2 and cavity C_1 is (21, 22)

$$\mathcal{H}_{\text{eff}} = -\frac{\hbar g^2}{\Delta}(a_1a_1^\dagger|c_2\rangle\langle c_2| - a_1^\dagger a_1|b_2\rangle\langle b_2|), \quad [12]$$

where g is a coupling coefficient, and a_1 and a_1^\dagger are destruction and creation operators for the field state inside cavity C_1 . The resulting entangled state between atom 2 and the field in cavity C_1 is $\frac{1}{2}(|0_1, a_2\rangle + |0_1, b_2\rangle + |1_1, a_2\rangle + e^{i\eta}|1_1, b_2\rangle)$, where $\eta = g^2\tau/\Delta_1$. The net result is that there is no photon number change inside the cavity, and there is a phase change only when there is one photon inside the cavity and the atom is in state




Fig. 2. (a) Atomic-level spacings for types *A* and *B* atoms. Here $\omega_a b_1 = \omega_c b_2 + \Delta_1$ and $\omega_a b_2 = \omega_c b_1 + \Delta_2$. (b) Schematics for the cavity quantum electrodynamics implementation of Grover's algorithm. Here all the atoms are initially in their ground states $|b\rangle$, i.e., $|b_1, b_2, b_3, b_4\rangle$. After passage through the box corresponding to the Walsh-Hadamard transformation \mathcal{W} , the state of the atoms is $(|a_1, a_2\rangle + |a_1, b_2\rangle + |b_1, a_2\rangle + |b_1, b_2\rangle)/2 \otimes |b_3, b_4\rangle$. After passage through box \mathcal{C} corresponding to the oracle for, say, state $|0, 1\rangle$, the atomic state is $(|b_1\rangle \otimes |a_3, a_2\rangle - |a_3, b_2\rangle + |b_3, a_2\rangle + |b_3, b_2\rangle)/2 \otimes |b_4\rangle$. Finally, after passage through box \mathcal{N} corresponding to the inversion about the average, the atomic state is $|b_1, b_2\rangle \otimes |a_3, b_4\rangle$.

$|b_2\rangle$. Interaction time τ and detuning Δ_1 are chosen such that $\eta = \pi$. This operation corresponds to the quantum phase gate discussed above and experimentally implemented in refs. 13 and 14.

Atom 3 of type *A* in ground state $|b_3\rangle$ now passes through cavity C_1 , followed by a classical field R_{C_1} . The interaction times with the cavity and classical fields are such that the cavity field is reduced to $|0\rangle$, and the resultant entangled state between atoms 2 and 3 is given by $(|a_3, a_2\rangle + |a_3, b_2\rangle + |b_3, a_2\rangle - |b_3, b_2\rangle)/2$. Here and in the following, we neglect the unimportant overall phase factor. Now $C_{0,0}$, $C_{0,1}$, $C_{1,0}$, and $C_{1,1}$ are implemented by turning classical fields R_3 and R_4 on, only R_3 on, only R_4 on, and none of the fields on, respectively. In the present example where the oracle picks $C_{0,1}$, field R_3 is tuned to frequency $\omega_a b_1$ and interacts with atom 3, and field R_4 tuned to frequency $\omega_a b_2$ is off. The interaction time for field R_3 is chosen such that $\theta = \pi/2$ and $\phi = 0$. Atoms 2 and 3 are in an entangled state $(|a_3, a_2\rangle - |a_3, b_2\rangle + |b_3, a_2\rangle + |b_3, b_2\rangle)/2$. This completes the oracle operation $C_{0,1}$ corresponding to the π phase shift of state $|a_3, b_2\rangle$.

Next, we implement operator \mathcal{N} , which inverts the states about the average. Atoms 3 and 2 interact with classical Ramsey field R_5 and R_6 , which are resonant with $|a\rangle$ to $|b\rangle$ transitions of atoms 3 and 2 with $\theta = \pi/4$ and $\phi = \pi/2$.

According to Eq. 12, we should next apply a quantum phase gate Q_α with $\alpha = \pi$. This step is accomplished as above by first transferring the atomic coherence of atom 2 to the empty cavity C_2 by adjusting the interaction time with the cavity appropriately. And then, on passing atom 3 through cavity C_2 , the entangled state of atom 3 and cavity C_2 is $(|a_3, 0_2\rangle - |a_3, 1_2\rangle + |b_3, 0_2\rangle - |b_3, 1_2\rangle)/2$.

Subsequently, a fourth atom of type *B* passes through cavity C_2 , and a classical field, R_{C_2} , leaves the cavity empty and transfers the field coherence to the atom yielding $(|a_3, a_4\rangle - |a_3, b_4\rangle + |b_3, a_4\rangle - |b_3, b_4\rangle)/2$. As a last step, atoms 3 and 4 pass through a sequence of classical fields, R_7 , resonant with $|a\rangle$ to $|b\rangle$ transitions of atom 3 and R_8 resonant with $|a\rangle$ to $|b\rangle$ transitions of atom 4, such that $\theta = \pi/4$, $\phi = \pi/2$ for R_7 , R_8 . This step completes the implementation of operator \mathcal{N} , and the resulting state of atoms 3 and 4 is $|a_3, b_4\rangle$. The final state of

atoms 3 and 4 would be $|a_3, a_4\rangle$, $|a_3, b_4\rangle$, $|b_3, a_4\rangle$ or $|b_3, b_4\rangle$, depending on the choice of $C_{\alpha,\beta}$.

Motivated by the above scheme, we consider an equivalent scheme that requires only one species of atoms and a single cavity, as shown in Fig. 3. Here the atoms have one lower level and two excited levels, whose splitting can be adjusted by applying electric or magnetic fields (23). These Stark or Zeeman splittings can be used to carry out resonant or dispersive atomic couplings with the cavity fields.

We consider a cavity that can support two modes of frequencies ν_1 and ν_2 . The three-level atoms having a lower level $|b\rangle$ and two upper levels $|a_1\rangle$ and $|a_2\rangle$ can have level spacings that depend on the applied electric or magnetic fields, as shown in Fig. 3a. Level $|b\rangle$ corresponds to qubit $|1\rangle$, and levels $|a_1\rangle$ and $|a_2\rangle$ correspond to qubit $|0\rangle$ for atoms 1 and 3 and atoms 2 and 4, respectively. In configuration 0, levels $|a_1\rangle$ and $|a_2\rangle$ are completely detuned with respect to cavity resonance frequencies ν_1 and ν_2 , and the atom is effectively decoupled from the cavity fields. In configuration 1, the atom resonantly interacts with ν_1 but is decoupled with ν_2 . In configuration 2, the atom interacts dispersively with ν_2 but is decoupled with ν_1 . Similarly, in configurations 3 and 4, the atom interacts resonantly with ν_2 with no interaction with ν_1 , dispersively with ν_1 , and with no interaction with ν_2 .

As before, four atoms are sent inside the cavity initially in their ground state $|b\rangle$ such that they interact with a sequence of classical driving fields R_i ($i = 1 - 8$) and cavity fields C_1 and C_2 of frequencies ν_1 and ν_2 , respectively, as shown in Fig. 3b. The interaction times are the same as in the first scheme. The appropriate interactions leading to the implementation of Grover's algorithm corresponding to the first scheme are obtained if the sequence of the atomic level spacings during the passage through R_i and C_i is chosen for various atoms as that given in Fig. 3c.

In conclusion, we have shown how Grover's algorithm can be implemented by using the known methods and techniques of quantum optics. Limitations are imposed by considerations of spontaneous emission and by cavity field damping. A potential source of error is the uncertainty associated with the location of atoms in the various fields.




Fig. 3. (a) Level spacings in three-level atoms corresponding to different applied electric or magnetic fields. (b) Applied classical fields R_i and cavity fields C_i on the incoming atoms. (c) Table indicating the configurations corresponding to Fig. 3a that the four atoms experience during their passage through various Ramsey and cavity fields.

We thank L. Davidovich, J. Dowling, K. Kapale, H. Lee, and C. P. Williams for valuable discussions. This research is supported by the

Office of Naval Research, the National Science Foundation, and the Welch Foundation.

1. Grover, L. K. (1997) *Phys. Rev. Lett.* **79**, 325–328.
2. Farhi, E. & Gutmann, S. (1998) *Phys. Rev. A* **57**, 2403–2406.
3. Lloyd, S. (2000) *Phys. Rev. A* **61**, 0103011–0103014.
4. Kwiat, P. G., Mitchell, J. R., Schwindt, P. D. D. & White, A. G. (1999) *J. Mod. Optics* **47**, 257–266.
5. Cerf, N. J., Adami, C. & Kwiat, P. G. (1998) *Phys. Rev. A* **57**, R1477–R1480.
6. Ahn, J., Weinacht, T. C. & Bucksbaum, P. H. (2000) *Science* **287**, 463–465.
7. Knight, P. (2000) *Science* **287**, 441–442.
8. Scully, M. O. & Zubairy, M. S. (2001) *Phys. Rev. A* **64**, 0223041–0223045.
9. Chuang, I. L., Gershenfeld, N. & Kubinec, M. (1998) *Phys. Rev. Lett.* **80**, 3408–3411.
10. Jones, J. A., Mosca, M. & Hansen, R. H. (1998) *Nature (London)* **393**, 344.
11. Jones, J. A. (1998) *Science* **280**, 229.
12. Braunstein, S. L., Caves, C. M., Jozsa, R., Linden, N., Popescu, S. & Schack, R. (1999) *Phys. Rev. Lett.* **83**, 1054–1057.
13. Rauschenbeutel, A., Nogues, G., Osnaghi, S., Bertet, P., Brune, M., Raimond, J. M. & Haroche, S. (1999) *Phys. Rev. Lett.* **83**, 5166–5169.
14. Turchette, Q. A., Hood, C. J., Lange, W., Mabuchi, H. & Kimble, H. J. (1995) *Phys. Rev. Lett.* **75**, 4710–4713.
15. Skarja, M., Borstnik, N. M., Loffler, M. & Walther, H. (1999) *Phys. Rev. A* **60**, 3229–3232.
16. Meschede, D., Walther, H. & Muller, G. (1985) *Phys. Rev. Lett.* **54**, 551–554.
17. Haroche, S. & Raimond, J. M. (1985) in *Advances in Atomic, Molecular, and Optical Physics*, eds. Bates, D. R. & Bederson, B. (Academic, New York), Vol. 20, p. 350.
18. An, K., Childs, J. J., Desari, R. R. & Feld, M. S. (1994) *Phys. Rev. Lett.* **73**, 3375–3378.
19. Raithel, G., Wagner, C., Walther, H., Narducci, L. M. & Scully, M. O. (1994) in *Advances in Atomic, Molecular, and Optical Physics*, ed. Berman, P. (Academic, New York), Suppl. 2, p. 57.
20. Hood, C. J., Chapman, M. S., Lynn, T. W. & Kimble, H. J. (1998) *Phys. Rev. Lett.* **80**, 4157–4160.
21. Brune, M., Haroche, S., Lefevre, V., Raimond, J. M. & Zagury, N. (1990) *Phys. Rev. Lett.* **65**, 976–979.
22. Scully, M. O. & Zubairy, M. S. (1997) *Quantum Optics* (Cambridge, London).
23. Maitre, X., Hagley, E., Nogues, G., Wunderlich, C., Goy, P., Brune, M., Raimond, J. M. & Haroche, S. (1997) *Phys. Rev. Lett.* **79**, 769–772.